Journal of Immunological Methods, 159 (1993) 293–294

© 1993 Elsevier Science Publishers B.V. All rights reserved 0022-1759/93/$06.00

Contents to volume 159 (1993)

Numbers 1–2 – 26 February 1993

Immunological standardization

The value of indirect immunofluorescence and solid phase techniques for ANCA detection. A report on the first phase of an international cooperative study on the standardization of ANCA assays
E.C. Hagen, K. Andrassy, E. Chernok, M.R. Daha, G. Gaskin, W. Gross, P. Lesavre, J. Lüdemann, C.D. Pusey, N. Rasmussen, C.O.S. Savage, A. Sinico, A. Wiik and F.J. Van der Woude EEC / BCR Group for ANCA Assay Standardization (Netherlands, France, Germany, Italy, Denmark, UK) 1

Research reports

Quantitative analysis of interleukin-2-induced proliferation in the presence of inhibitors using a mathematical model
E.V. Svirshchevskaya, I.A. Sidorov, N.Y. Viskova and I.M. Dozmorov (Russian Federation) 17

Autologous lymphocytes prevent the death of monocytes in culture and promote, as do GM-CSF, IL-3 and M-CSF, their differentiation into macrophages
M. Lopez, C. Martinache, S. Canepa, M. Chokri, F. Scotto and J. Bartholeyns (France) 29

Use of particulate forms of protein antigens to increase the sensitivity of antigen-specific proliferation assays
J.-Y. Wu, K.J. Munroe, B.H. Gardner, M.S. Wyand and M.J. Newman (USA) 39

Use of logistic regression in comparing multiple limiting dilution analyses of antigen-reactive T cells
J.A. Burleson, T.A. Binder and I. Goldschneider (USA) 47

Quantitative immunoassays for type II collagen and its cyanogen bromide peptides
G.R. Srinivas, H.J. Barrach and C.O. Chichester (USA) 53

Isolation of a protein complex from purulent sputum consisting of proteinase-3 and α1-antitrypsin reactive with anti neutrophil cytoplasmic antibodies
B.E.P.B. Ballieux, E.C. Hagen, C. Van der Keur, N.D. Zegers, L.A. Van Es, F.J. Van der Woude and M.R. Daha (Netherlands) 63

A microneutralization enzyme immunoassay for antibody to human cytomegalovirus
T. Tomiyama, T. Sugano, S. Tani, K. Hosoda and Y.-i. Matsumoto (Japan) 71

Optimization of the L-M cell bioassay for quantitating tumor necrosis factor α in serum and plasma
F.E. Nargi and T.J. Yang (USA) 81

Rapid analysis of leukocyte-endothelial adhesion
A.A. Vaporciyan, M.L. Jones and P.A. Ward (USA) 93

Liposome immunoassay (LIA) with antigen-coupled liposomes containing alkaline phosphatase
C.-K. Kim and S.-J. Lim (South Korea) 101

A new cell culture method (the lateral diffusion system) suitable for the induction of antibody-forming cells in vitro
F.M. Ubeira, P. Baltar and F. Romarís (Spain) 107

Isolation of cytotoxic Kupffer cells by a modified enzymatic assay: a methodological study
G. Heuff, J.J.E. Steenbergen, A.A. Van de Loosdrecht, I. Sirovich, C.D. Dijkstra, S. Meyer and R.H.J. Beelen (Netherlands) 115

Purification of monoclonal antibodies from tissue culture medium depleted of IgG
C.R. Darby, K. Hamano and K.J. Wood (UK) 125

An automated micro-fluorometric assay for monitoring oxidative burst activity of phagocytes
C.P. Wan, E. Myung and B.H.S. Lau (USA) 131

Efficient preparation of human monoclonal antibody-secreting heterohybridomas using peripheral B lymphocytes cultured in the CD40 system
A. Darveau, M.-C. Chevrier, S. Néron, R. Delage and R. Lemieux (Canada) 139

A simple optical fiber device for quantitative fluorescence microscopy of single living cells
M. Van Graft, B. Oosterhuis, K.O. Van der Werf, B.G. De Grooth and J. Greve (Netherlands) 145

Application of polymer β-cyclodextrin-treated fetal calf serum to in vitro antibody production
I. Yamamoto, E. Miyai and T. Chiba (Japan) 153

A multiparameter flow cytometric method to study surface molecules involved in interactions between subpopulations of cells
J.-P. Aubry, J.G. Shields, K.U. Jansen and J.-Y. Bonnefoy (Switzerland) 161
Simultaneous evaluation of phagocytosis and FcγR-mediated oxidative burst in human monocytes by a simple flow cytometry method
J.A. Casado, J. Merino, J. Cid, M.L. Subira and A. Sanchez-Ibarrola (Spain) 173
Covalent linking of haptens, proteins and nucleic acids to a modified polystyrene support
A. Niveleau, D. Sage, C. Reynaud, C. Bruno, S. Legastelois, V. Thomas and R. Dante (France) 177
Optimizing follicular dendritic cell isolation by discontinuous gradient centrifugation and use of the magnetic cell sorter (MACS)
J. Schmitz, S. Petrasch, J. Van Lunzen, P. Racz, H.-D. Kleine, F. Hufert, P. Kern, H. Schmitz and K. Tenner-Racz (Germany) 189
Detection of intracellular cytokines by flow cytometry
T. Jung, U. Schauer, C. Heusser, C. Neumann and C. Rieger (Germany, Switzerland) 197
In vitro phagocytosis inhibition assay for the screening of potential candidate antigens for sub-unit vaccines against the asexual blood stage of *Plasmodium falciparum*
J. Gysin, S. Gavoille, D. Mattei, A. Scherf, S. Bonnefoy, O. Mercereau-Puijalon, T. Feldmann, B. Müller-Hill and L. Pereira da Silva (French Guiana, France, Germany) 209
A novel ELISA assay for the detection of C3 nephritic factor
J. Seino, A.W.L. V.d. Wall Bake, L.A. Van Es and M.R. Daha (Netherlands) 221
Nanogram quantitation of secreted protein in a recombinant yeast fermentation using an immuno-ligand assay
R.S.R. Robinett and W.K. Herber (USA) 229
Quantitative measurement of human cytokine gene expression by polymerase chain reaction
P.C. Melby, B.J. Darnell and V.V. Tryon (USA) 235
Serum antibody production by oral immunization with dextran B512
M. Tanaka, Y. Takeda and S. Takeo (Japan) 245
A colorimetric assay to evaluate the immune reactivity of blood platelets based on the reduction of a tetrazolium salt
D. Vanhée, M. Joseph, H. Vorng and A.-B. Tonnel (France) 253
Immunochemical characterization of anti-H monoclonal antibodies obtained from a mouse immunized with human saliva
T. Nakajima, S. Yazawa, S. Miyazaki and K. Furukawa (Japan) 261
The use of recombinant human growth hormone for radioiodination and standard preparation in radioimmunoassay
M.T.C.P. Ribela, Y. Murata, L. Morganti, D. Toniolo and P. Bartolini (Brazil, Italy) 269

Technical note
Splenic implantation of nitrocellulose pellets as a means of raising antibodies to nanogram quantities of antigen
J.M.S. Forrest and H.A. Ross (UK) 275

Short communications
Measurement of antibody binding to antigenic peptides conjugated in situ to albumin-coated microtitre plates
M.R. Price, M. Sekowski, D.S.W. Hooi, L.G. Durrant, F. Hudecz and S.J.B. Tendler (UK, Hungary) 277
Use of an enzyme-linked immunosorbent assay (ELISA) to measure the expression of the K88 fimbrial antigen by enterotoxigenic *Escherichia coli* (ETEC)
D. Payne, N. Moore and P. Lambert (UK) 283

Letter to the editors
Long term preservation of indirect immunofluorescent specimens in ultralow freezer (−75°C)
Y. Wang and K. Ben (People's Republic of China) 291

Contents to volume 159 (1993) 293

Notes to authors 295